

RESEARCH

Open Access



Sleep disturbances and predictors of nondeployability among active-duty Army soldiers: an odds ratio analysis of medical healthcare data from fiscal year 2018

Jaime K. Devine^{1*}, Jacob Collen², Jake J. Choynowski³ and Vincent Capaldi³

Abstract

Background: The impact of sleep disorders on active-duty soldiers' medical readiness is not currently quantified. Patient data generated at military treatment facilities can be accessed to create research reports and thus can be used to estimate the prevalence of sleep disturbances and the role of sleep on overall health in service members. The current study aimed to quantify sleep-related health issues and their impact on health and nondeployability through the analysis of U.S. military healthcare records from fiscal year 2018 (FY2018).

Methods: Medical diagnosis information and deployability profiles (e-Profiles) were queried for all active-duty U.S. Army patients with a concurrent sleep disorder diagnosis receiving medical care within FY2018. Nondeployability was predicted from medical reasons for having an e-Profile (categorized as sleep, behavioral health, musculoskeletal, cardiometabolic, injury, or accident) using binomial logistic regression. Sleep e-Profiles were investigated as a moderator between other e-Profile categories and nondeployability.

Results: Out of 582,031 soldiers, 48.4% ($n = 281,738$) had a sleep-related diagnosis in their healthcare records, 9.7% ($n = 56,247$) of soldiers had e-Profiles, and 1.9% ($n = 10,885$) had a sleep e-Profile. Soldiers with sleep e-Profiles were more likely to have had a motor vehicle accident (pOR (prevalence odds ratio) = 4.7, 95% CI 2.63–8.39, $P \leq 0.001$) or work/duty-related injury (pOR = 1.6, 95% CI 1.32–1.94, $P \leq 0.001$). The likelihood of nondeployability was greater in soldiers with a sleep e-Profile and a musculoskeletal e-Profile (pOR = 4.25, 95% CI 3.75–4.81, $P \leq 0.001$) or work/duty-related injury (pOR = 2.62, 95% CI 1.63–4.21, $P \leq 0.001$).

Conclusion: Nearly half of soldiers had a sleep disorder or sleep-related medical diagnosis in 2018, but their sleep problems are largely not profiled as limitations to medical readiness. Musculoskeletal issues and physical injury predict nondeployability, and nondeployability is more likely to occur in soldiers who have sleep e-Profiles in addition to these issues. Addressing sleep problems may prevent accidents and injuries that could render a soldier nondeployable.

Keywords: Medical readiness, Behavioral sleep medicine, Deployability, Healthcare records, Military, Big data, Data mining

* Correspondence: jdevine@ibrinc.org

¹Institutes for Behavior Resources, Operational Fatigue and Performance, 2104 Maryland Ave, Baltimore, MD 21218, USA

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Readiness is the number one priority in the U.S. Army [1]. The Army's ability to restore and regenerate equipment, expertise and personnel are key to its ongoing success [2]. Soldiers who are unable to deploy when the Army needs them directly affect this readiness. Reducing the rate of nondeployability among soldiers is a complex challenge that Army organizations and senior staff need to address to maximize readiness [3].

Decreasing the number of nondeployable soldiers across the Army depends on determining and mitigating the underlying causes for nondeployable conditions. Soldiers can be categorized as nondeployable due to administrative, legal or medical conditions. Notably, medically nondeployable soldiers constitute the largest category, accounting for approximately 80% of nondeployable soldiers [3, 4]. Medical readiness, therefore, is an important target for reducing nondeployability in the Army.

Understanding which medical conditions most greatly contribute to nondeployability is essential in developing strategies for preventative care and risk mitigation. The Army maintains a centralized authoritative database of medical readiness information of Army personnel known as the Medical Occupational Data System (MODS). Within the MODS are two Web-based modules that track and record medical readiness information. The Medical Protection System (MEDPROS) is the primary tool to record, track, and report soldiers' medical conditions, and the electronic profiling system (e-Profile) tracks whether any medical conditions may render soldiers medically unable to deploy on a temporary or permanent basis [5, 6]. Soldiers can acquire a "profile" in six different categories: physical functional capacity (P), upper extremities (U), lower extremities (L), hearing and ears (H), eyes and vision (E), and psychiatric (S). Together, the six categories are often referred to as "PULHES". Soldiers with e-Profiles have a score between 1 and 4 for each PULHES category. A score of 1 indicates that the soldier is medically sound in that category, while a score of 2 indicates a mild impairment (i.e., a soldier with an E (eyes and vision) score of 2 may require glasses). A permanent e-Profile score of 3 or higher in any PULHES category indicates that the soldier is medically undeployable. For example, an E score of 4 may indicate blindness. Soldiers can have multiple concurrent e-Profiles for multiple medical conditions.

Sleep is a factor related to health concerns across the medical spectrum. Sleep disorders in and of themselves are debilitating [7, 8], but sleep is also related to physical and mental health [9–12]. For example, individuals with insomnia or obstructive sleep apnea (OSA) are at greater risk for obesity, type 2 diabetes, atherosclerosis, coronary heart disease, heart failure, hypertension, stroke, and trauma-related nightmares [9, 13]. Moreover, fatigue is a contributing factor to the occurrence of traffic and

workplace accidents [14–17]. It is therefore possible that sleep problems could contribute to medical conditions that result in permanent e-Profiles. However, while sleep complaints and sleep disorders are common among soldiers [13, 18–21], the impact of sleep-related health issues on deployability within active-duty service members is not currently quantified.


Understanding the scope of sleep problems and their impact on deployability will help command leadership and medical professionals determine where to concentrate efforts to treat and prevent debilitating health issues among service members and maintain the medical readiness of the U.S. Army. The current study aimed to quantify the prevalence of sleep-related health issues through the analysis of U.S. military healthcare records from fiscal year 2018 (FY2018) to assess the relationship between sleep disorders and health and nondeployability in active-duty soldiers.

Methods

Patients data generated through the MODS, MEDPROS and e-Profile are stored in the Composite Healthcare System (CHCS). The Standard Inpatient Data Record (SIDR) is extracted from the CHCS database twice per month and transmitted securely to the Patient Administration Systems and Biostatistics Activity (PASBA). The Comprehensive Ambulatory/Professional Encounter Record (CAPER) is extracted from the CHCS database daily and transmitted securely to PASBA. The data in the PASBA can be accessed by submitting a data extraction request and data sharing agreement to the Defense Health Agency (DHA) and the Office of the Surgeon General (OTSG) for research and for decision support activities to enable the U.S. Army Medical Command to operate effectively.

Data for the current study were requested from, de-identified and compiled by a PASBA DHA health statistician prior to delivery/analysis, as depicted in Fig. 1. Medical diagnosis information and e-Profile status data were queried for all active-duty U.S. Army patients with a concurrent sleep disorder diagnosis receiving medical care within fiscal year 2018. Sleep disorders were queried using International Statistical Classification of Diseases and Related Health Problems, version 10 (ICD-10) medical diagnosis codes in the following ranges: G47.00-G47.9: Sleep disorder; F51.01-F51.9: Sleep disorders not due to a substance or known physiological condition; and Z72.820-Z72.821: Problems related to sleep. A fictional linking identifier was created to replace Department of Defense medical identifiers for all patient information. Data were compiled using Excel 2013 (Microsoft Corp., Redmond, WA, USA) and Access 2013 (Microsoft Corp., Redmond, WA, USA).

SPSS statistical software (version 25, SPSS, Inc., Chicago, IL, USA) was used to conduct statistical analysis of de-



identified e-Profile data. An alpha level of 0.05 was used for all statistical tests. Binary variables were calculated from the e-Profile classification system called “reason for visit” such that sleep e-Profiles were defined as “reason for visit = sleep or focus area: sleep”, behavioral health e-Profiles were defined as “reason for visit = behavioral health” (with the exception of sleep disorders, which were categorized under sleep e-Profiles) and cardiometabolic health e-Profiles were defined as “reason for visit = cardiology, endocrine/general, or gastroenterology”. Soldiers with any e-Profile (temporary or permanent, all scores) due to that condition were categorized as having the condition (coded as 1), while soldiers with only e-Profiles for any other medical condition were categorized as not having the condition (coded as 0).

Binary variables for all causes of nondeployability, nondeployability due to a sleep disorder, nondeployability due to behavioral health, and nondeployability due to cardiometabolic health were calculated from PULHES scores such that all temporary e-Profiles and permanent e-Profiles with scores ≤ 2 in all categories were considered deployable and a permanent e-Profile PULHES score ≥ 3 was considered nondeployable. Additionally, medical records due to motor vehicle accidents (defined as injury mechanism: traffic accident or motor vehicle accident) and work/duty-related injuries (defined as injury mechanism: battle injury, duty-related injury, or work-/task-related injury) were used to create binary variables for e-Profiles due to accident or work/duty-related injury.

Chi-squared tests for independence examined the relationship between e-Profiles for sleep, behavioral health, cardiometabolic health, work-related injuries and accidents and deployability. *P* values were considered significant when *p* was equal to or less than 0.05. Binomial logistic regressions predicted the likelihood of nondeployability for soldiers who had an e-Profile due to sleep, behavioral health, cardiometabolic health, work-related injury or an accident compared to those who did not, as well as the likelihood of having an e-Profile for behavioral health, cardiometabolic health, work/duty-related injury or motor vehicle accident for soldiers with or without an e-Profile for sleep using block entry [22]. Further, interaction effects between sleep e-Profiles and other variables of interest (behavioral health, cardiometabolic health, work/duty-related injury or motor vehicle accidents) were calculated to examine moderation of the relationship between predictors of nondeployability and sleep e-Profiles.

Results

Soldier and e-profile descriptives

There were on average 582,031 active-duty Army soldiers in FY2018. Of this active-duty population, 48.4% ($n = 281,738$) soldiers fit the search criteria of having sought medical treatment for a sleep-related condition (as defined by ICD-10 codes) in FY2018 and 9.7% ($n = 56,247$) had a concurrent e-Profile tracking a temporary or permanent medical conditions that may have rendered them medically not ready to deploy. Soldiers

with e-Profiles ranged from 17 to 66 years old (mean ± SD: 36.5 ± 9.4; median: 37) and the average body mass index (BMI) was 29.0 ± 4.1 kg/m². The majority of soldiers with e-Profiles were male (71.3%, *n* = 40,080).

About 99 % (99.7%, *n* = 56,082) of soldiers had seven e-Profiles or less. The remaining 0.3% (*n* = 165) of soldiers had between 8 and 20 e-Profiles. Musculoskeletal conditions (such as injury to the spine, bones or joints) were the most commonly listed condition for e-Profiles, accounting for 71.8% (*n* = 40,508) of all e-Profiles. Sleep disorders were the listed condition for 19.4% (*n* = 10,885) of all e-Profiles from soldiers with sleep-related diagnoses in FY2018. Specific diagnoses associated with sleep e-Profiles are summarized in Table 1.

About 16 % of soldiers with e-Profiles (16.2%, *n* = 9119) were classified as nondeployable. The leading reason for nondeployability was musculoskeletal conditions, accounting for 47.9% (*n* = 4366) of nondeployability. Behavioral health was the listed reason for nondeployability in 27.2% (*n* = 2481) of nondeployable soldiers. Cardiometabolic health was the listed reason for nondeployability in 8.1% (*n* = 736) of the nondeployable soldiers. Sleep disorders were the listed reason for nondeployability in 1.8% (*n* = 165) of nondeployable soldiers. The listed reasons for nondeployability in the remaining 15.0% (*n* = 1371) of soldiers were related to a wide range of health conditions, such as urological, pulmonary, and degenerative conditions.

Prevalence odds ratios for comorbid conditions in relation to sleep e-profiles

The comorbidity of sleep e-Profiles and musculoskeletal e-Profiles, cardiometabolic e-Profiles, and behavioral health e-Profiles, as well as the prevalence of motor vehicle accidents, work/duty-related injuries and nondeployability, are summarized in Table 2. Comorbidities of sleep profiles were statistically significant for all e-Profile types (*P* < 0.05)

Table 1 Sleep disorder diagnoses in soldiers with e-Profiles

Listed condition	Case	Percentage of soldiers with sleep e-Profile (%)	Percentage of soldiers with Sleep-related diagnosis (%)	Percentage of active-duty population (%)
Sleep e-Profile (All)	10,885	100.0	19.4	1.9
Obstructive sleep apnea e-Profiles	10,442	96.0	3.7	1.8
Narcolepsy and circadian disorder e-Profile	190	1.7	0.07	0.03
Insomnia e-Profile	107	1.0	0.04	0.02
Narcolepsy and OSA e-Profiles	30	0.3	0.01	0.005

Breakdown of sleep e-Profiles by listed condition for active-duty soldier populations from FY2018. OSA was the listed condition for the majority of soldiers with sleep e-Profiles and represented 1.9% of all active-duty soldiers

Table 2 Prevalence odds ratios of having a sleep e-Profile and other e-Profile category, accident, injury, and nondeployability

Item	Sleep e-Profile [<i>n</i> (%)]		pOR	95%CI	P-Value
	Yes	No			
Nondeployable					
Yes	2027(3.6)	7092(12.6)	0.81	0.77–0.85	<i>P</i> ≤ 0.001
No	8858(15.8)	38,270(68.0)			
Musculoskeletal e-Profile					
Yes	6740(12.0)	34,470(61.2)	0.51	0.49–0.54	<i>P</i> ≤ 0.001
No	4145(7.4)	10,892(19.4)			
Cardiometabolic e-Profile					
Yes	46(0.1)	1322(2.5)	0.67	0.60–0.75	<i>P</i> ≤ 0.001
No	10,418(19.0)	44,040(78.4)			
Behavioral health e-Profile					
Yes	1040(1.8)	4470(8.0)	1.04	0.96–1.11	<i>P</i> = 0.34
No	9845(17.5)	40,892(72.7)			
Motor vehicle accident					
Yes	12(0.1)	234(0.4)	4.7	2.63–8.39	<i>P</i> ≤ 0.001
No	10,873(19.3)	45,128(80.2)			
Work/duty-related injury					
Yes	120(0.2)	794(1.4)	1.6	1.32–1.94	<i>P</i> ≤ 0.001
No	10,765(19.1)	44,568(79.3)			

Prevalence odds ratio analysis of the likelihood of comorbidity between sleep e-Profiles and other e-Profile types. Soldiers with sleep e-Profiles were more likely to have a musculoskeletal e-Profile, motor vehicle accident, or work/duty-related injury than soldiers without a sleep e-Profile

except for behavioral health (*P* = 0.34). Soldiers with sleep e-Profiles were more likely to also have had a motor vehicle accident or work/duty-related injury compared to soldiers without a sleep e-Profile.

Prevalence odds ratios for nondeployability in relation to having a musculoskeletal, cardiometabolic or behavioral health e-Profile, motor vehicle accident or work/duty-related injury are summarized in Table 3. Nondeployability was significantly related to all e-Profile types (all *P* ≤ 0.001; data not shown). Nondeployable soldiers were significantly more likely to have musculoskeletal e-Profiles or have had a work/duty-related injury. In addition, there was also a trend (*P* = 0.06) for nondeployable soldiers to have had a motor vehicle accident.

The moderation effects of having a sleep e-Profile on the odds ratios for nondeployability were examined for musculoskeletal e-Profiles, cardiometabolic e-Profiles, behavioral health e-Profiles, motor vehicle accidents and work/duty-related injuries. Table 4 summarizes the interaction between sleep e-Profiles and the three variables found to predict greater odds of nondeployability (musculoskeletal e-Profiles, motor vehicle accidents and work/duty-related injuries). Soldiers with a musculoskeletal e-Profile and a sleep e-Profile were significantly more likely to be nondeployable, and soldiers with a work/duty-

Table 3 Prevalence odds ratios of nondeployability and profile categories, motor vehicle accidents and work/duty-related injuries

Item	Nondeployable [n(%)]		pOR	95%CI	P-value
	Yes	No			
Musculoskeletal profile					
Yes	6174(11.0)	34,234(60.9)	1.27	1.21–1.33	$P \leq 0.001$
No	2945(5.2)	12,894(22.9)			
Cardiometabolic profile					
Yes	851(1.5)	938(1.7)	0.2	0.18–0.22	$P \leq 0.001$
No	8268(14.7)	46,190(82.1)			
Behavioral health profile					
Yes	2894(5.1)	2616(4.6)	0.13	0.12–0.13	$P \leq 0.001$
No	6225(11.1)	44,512(79.1)			
Motor vehicle accident					
Yes	29(0.1)	217(0.4)	1.45	0.98–2.14	$P = 0.06$
No	9090(16.1)	46,911(83.4)			
Work/duty-related injury					
Yes	109(0.2)	805(1.4)	1.44	1.17–1.76	$P \leq 0.001$
No	9010(16.0)	46,323(82.4)			

Prevalence odds ratio analysis of the likelihood of being nondeployable and having a musculoskeletal, cardiometabolic or behavioral health e-Profile, motor vehicle accident or work/duty-related injury. Nondeployable Soldiers were significantly more likely to have musculoskeletal e-Profiles or have had a work/duty-related injury

related injury and sleep e-Profile were nearly three times as likely to be nondeployable. The odds ratio of nondeployability given that a soldier had an incidence of a motor vehicle accident in addition to a sleep e-Profile was not statistically significant.

Discussion

Nearly half of active-duty U.S. Army soldiers had a diagnosis for a sleep-related issue during fiscal year 2018. However, only 3.8% of these soldiers additionally had an

e-Profile for their sleep condition. This may indicate that sleep disorders are not generally considered by military clinicians to impact medical readiness to a degree sufficient to warrant the generation of an e-Profile. Even when profiled, soldiers with a sleep, cardiometabolic, or behavioral health e-Profile were less likely to be nondeployable. These statistics indicate that sleep, cardiometabolic and behavioral health disorders do not greatly influence the medical readiness and deployability of U.S. Army soldiers.

In contrast, musculoskeletal issues accounted for 47.9% of the nondeployability of soldiers. Soldiers who had a motor vehicle accident or work/duty-related injury were also more likely to be nondeployable (Table 3). Interestingly, soldiers with sleep e-Profiles were over one and a half times more likely to also experience accidents or injuries as soldiers with no sleep e-Profile.

Subsequently, we examined the moderation of predictors of nondeployability by comorbid sleep e-Profiles. Soldiers with a musculoskeletal e-Profile or work/duty-related injury in addition to a sleep e-Profile were between two to four times more likely to be nondeployable as soldiers with either one or none of these issues (Table 4). The odds ratio for having a both motor vehicle accident and sleep e-Profile and being nondeployable was not significant, most likely due to the small sample size of soldiers with an e-Profile for motor vehicle accidents ($n = 246$). While it is impossible to determine causal relationships between predictors of nondeployability and sleep from these data, the picture begins to emerge that sleep disorders are connected to medical readiness in indirect and undocumented ways, such as through correlation with injuries.

Interestingly, having a sleep e-Profile was not related to an increased likelihood of having a behavioral health e-Profile. This finding contradicts the known relationship between mental health and sleep from the literature

Table 4 Prevalence odds ratios of nondeployability, predictors of nondeployability and sleep e-Profiles

Item	Nondeployable [n(%)]		pOR	95%CI	P-value
	Yes	No			
Musculoskeletal e-Profile and Sleep e-Profile					
Yes	1565(2.8)	5116(9.1)	4.25	3.75–4.81	$P \leq 0.001$
No	7554(13.4)	42,012(74.7)			
Motor Vehicle Accident and Sleep e-Profile					
Yes	3(0.0)	9(0.0)	2.16	0.55–8.52	$P = 0.27$
No	9116(16.2)	47,119(83.8)			
Work/Duty-Related Injury and Sleep e-Profile					
Yes	31(0.1)	89(0.2)	2.62	1.63–4.21	$P \leq 0.001$
No	9088(16.1)	47,039(83.6)			

Prevalence odds ratio analysis comparing the likelihood of having a sleep e-Profile in combination with either a musculoskeletal e-Profile, motor vehicle accident or work/duty-related injuries. Soldiers with a musculoskeletal e-Profile and a sleep e-Profile or a work/duty-related injury and a sleep e-Profile were more likely to be nondeployable than soldiers with only one or neither of those conditions

[10, 19, 23]. Additionally, soldiers with sleep e-Profiles were less likely to have a cardiometabolic profile (Table 2). The current data pertain to medical diagnoses or medical readiness information as determined by a military clinician. Many cardiometabolic and behavioral health disorders are comorbid with sleep disorders or have sleep disturbances as a symptom [9, 19, 24–27]. This overlap in symptomology may not translate to having an e-Profile when clinicians are asked to categorize a soldier's limitations to medical readiness using the PULHES scoring system. Additionally, considering that soldiers with e-Profiles for sleep, cardiometabolic or behavioral health issues were less likely to be nondeployable, it is possible that these conditions are not generally considered serious enough to limit a soldier's ability to serve. For example, OSA, which accounted for the vast majority of sleep disorder diagnoses in the current study, can be treated with oral devices, surgery or positive airway pressure devices [28] and thus would not necessarily limit a soldier's functionality.

Additionally, an underappreciation for the severity of sleep problems may prevent soldiers from discussing fatigue-related issues with their healthcare providers. While the Department of Defense recognizes sleep as an important component of health and the performance triad [29], Army culture has not traditionally shared this respect [30]. Undervaluing the need for sleep likely contributes to the prevalence of sleep disturbances in active-duty soldiers through poor sleep hygiene but may also mask the severity of sleep problems in soldiers who downplay the importance of rest or who think that complaining about sleep may be interpreted as a sign of weakness.

It should be noted that the e-Profile data extracted for these analyses were only from active-duty soldiers with a sleep-related diagnosis in fiscal year 2018. Therefore, even though the sleep disorders accounted for e-Profiles in only 19.4% of soldiers with e-Profiles, the entire population ($n = 281,738$) from these analyses suffered from some sleep-related health issue in 2018. It is therefore unclear whether the findings from the current analyses can be generalized to the entire active-duty Army population or whether only motor vehicle accidents, work/duty-related injuries and musculoskeletal issues are predictors of nondeployability in soldiers with sleep problems. Future analyses will compare soldiers with sleep disorders to soldiers with no sleep complaints.

The current analyses are limited not only by the population but also by the source of the data. Medical healthcare data are a largely untapped resource for examining the complex manifestation of health issues, but the MODS, MEDPRO, e-Profile, SIDR or CAPER merely track medical encounters; they are not designed for hypothesis testing. As data mining techniques and the field of data science advance, there are bound to be improvements in the

modeling of medical healthcare data for research needs. The interpretation of findings is limited by the disproportionate numbers of e-Profiles due to different causes, which may be indicative of actual prevalence but is less than ideal for hypothesis testing, such as the high prevalence of musculoskeletal e-Profiles ($n = 40,508$) versus the relative rarity of e-Profiles due to motor vehicle accident ($n = 246$). Moreover, the current analyses should be considered correlative rather than causative.

Despite limitations to the analyses and interpretation, the data suggest that sleep disorders are a prodigious issue in the U.S. Army. A soldier is 4.7 times more likely to have a motor vehicle accident if he or she also has a profiled sleep disorder as a soldier with no sleep e-Profile. Importantly, a soldier was more likely to be nondeployable if he or she jointly had a musculoskeletal e-Profile and a sleep e-Profile than if he or she had only one of the two e-Profiles. These numbers highlight the impact of sleep on seemingly unrelated medical issues. A recent study by Shattuck et al. likewise showed an association between musculoskeletal complaints and shorter nighttime sleep as well as an increased report of fatigue in crewmembers on a U.S. Navy aircraft carrier [31], indicating that the comorbidity of sleep and musculoskeletal issues may be an epidemic across the Armed Services.

Conclusions

In addition to half of active-duty soldiers being diagnosed with sleep-related issues, sleep seems to be negatively impacting the medical readiness of service members. Musculoskeletal issues and physical injury predict nondeployability, and the occurrence of these problems is related to having a sleep disorder. Sleep disturbances constitute an underlying risk to medical readiness. Decreasing risk to active-duty military should go beyond the obvious causes and address underlying issues such as sleep disturbances to improve medical readiness and maximize the health of all U.S. service members.

Abbreviations

BMI: Body mass index; CAPER: Comprehensive ambulatory/professional encounter record; CHCS: Composite healthcare system; DHA: Defense health agency; E-Profile: Electronic profiling system; FY2018: Fiscal year 2018; ICD-10: International statistical classification of diseases and related health problems, version 10; MEDPROS: Medical protection system; MODS: Medical occupational data system; OSA: Obstructive sleep apnea; OTSG: Office of the surgeon general; PASBA: Patient administration systems and biostatistics activity; POR: Prevalence odds ratio; PULHES: Physical functional capacity (P), upper extremities (U), lower extremities (L), hearing and ears (H), eyes and vision (E), and psychiatric (S); SIDR: Standard Inpatient Data Record.

Acknowledgments

This material has been reviewed by the Walter Reed Army Institute of Research. There is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the true views of the Department of the Army or the Department of Defense. The authors would like to thank the DHA health statisticians Debra Greco and Armen Zakharyan for their help in extracting the requested data.

Authors' contributions

JKD analyzed and interpreted all data presented in this paper and served as the primary author for the manuscript. JJC compiled all data received from the MEDPROS, e-Profile, or other healthcare sources into one cohesive database and managed the data extractions from PASBA. JC and VC provided expertise on military healthcare, sleep medicine, and accessing the Medical Occupational Data System and were contributing authors. All authors read and approved the final manuscript.

Funding

The Department of Defense Military Operational Medicine Research Program (MOMRP) supported this study.

Availability of data and materials

The datasets generated and/or analyzed during the current study are not publicly available due to the potential sensitive nature of military healthcare data.

Ethics approval and consent to participate

This protocol was deemed exempt because it did not involve human participants but utilized only de-identified human medical record data. All study procedures were approved by the Walter Reed Army Institute of Research Institutional Review Board (DoD Assurance: A20126; HHS Federal Wide Assurance: #0000015; IRB Registration: #00000794).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Institutes for Behavior Resources, Operational Fatigue and Performance, 2104 Maryland Ave, Baltimore, MD 21218, USA. ²Pulmonary, Critical Care and Sleep Medicine Walter Reed National Military Medical Center, Bethesda, MD 20889, USA. ³Behavioral Biology Branch, Center for Military Psychiatry and Neuroscience, Walter Reed Army Institute of Research, Silver Spring, MD 20910, USA.

Received: 23 July 2019 Accepted: 26 February 2020

Published online: 10 March 2020

References

- Milley TM. Readiness for ground combat is No. 1 priority. 2015. <https://www.armytimes.com/news/pentagon-congress/2015/08/28/milley-readiness-for-ground-combat-is-no-1-priority/>. Accessed 19 May 2019.
- Nataraj S, Markel MW, Hastings JL, Larson EV, Luoto JE, Maerzluff CE, et al. Evaluating the Army's ability to regenerate: history and future options. Santa Monica: RAND Corporation; 2018. <https://apps.dtic.mil/dtic/tr/fulltext/u2/105515.pdf>. ISBN: 978-0-8330-9663-0.
- Arnold S, Crate C, Drennan S, Gaylord J, Hoffmann A, Martin D, et al. Non-deployable soldiers: understanding the Army's challenge. Carlisle Barracks: US Army War College; 2011. <https://apps.dtic.mil/dtic/tr/fulltext/u2/a560651.pdf>.
- Cox M. Army releases deploy-or-out rules for administratively sidelined troops. 2018. <https://www.military.com/daily-news/2018/11/13/army-releases-deploy-or-out-rules-administratively-sidelined-troops.html>. Accessed 26 August 2019.
- Williams A. MEDPROS an important part of Soldiers' readiness. 2012. https://www.army.mil/article/83121/medpros_an_important_part_of_soldiers_readiness. Accessed 20 May 2019.
- Stiltner J. Why is MEDPROS important? 2012. https://www.army.mil/article/76221/why_is_medpros_important. Accessed 21 May 2019.
- Luyster FS, Strollo PJ Jr, Zee PC, Walsh JK. Boards of directors of the American academy of sleep medicine, the sleep research society. sleep: a health imperative. *Sleep*. 2012;35(6):727–34.
- Grandner MA. Sleep, health, and society. *Sleep Med Clin*. 2017;12(1):1–22.
- St-Onge MP, Grandner MA, Brown D, Conroy MB, Jean-Louis G, Coons M, et al. Sleep duration and quality: impact on lifestyle behaviors and cardiometabolic health: a scientific statement from the American heart association. *Circulation*. 2016;134(18):e367–86.
- Devine JK, Wolf JM. Integrating nap and night-time sleep into sleep patterns reveals differential links to health-relevant outcomes. *J Sleep Res*. 2016;25(2):225–33.
- Ritland BM, Simonelli G, Gentili RJ, Smith JC, He X, Oh H, et al. Sleep health and its association with performance and motivation in tactical athletes enrolled in the reserve Officers' training corps. *Sleep Health*. 2019;5(3):309–14.
- Palmer CA, Alfano CA. Sleep and emotion regulation: an organizing, integrative review. *Sleep Med Rev*. 2017;31:6–16.
- Cremer JL, Brock MS, Mysliwiec V. Nightmares in United States military personnel are multifactorial and require further study. *J Clin Sleep Med*. 2018;14(7):1275–6.
- Zhang G, Yau KK, Zhang X, Li Y. Traffic accidents involving fatigue driving and their extent of casualties. *Accid Anal Prev*. 2016;87:34–42.
- Uehli K, Mehta AJ, Miedinger D, Hug K, Schindler C, Holsboer-Trachsler E, et al. Sleep problems and work injuries: a systematic review and meta-analysis. *Sleep Med Rev*. 2014;18(1):61–73.
- Akerstedt T, Wright KP Jr. Sleep loss and fatigue in shift work and shift work disorder. *Sleep Med Clin*. 2009;4(2):257–71.
- Sparrow AR, Mollicone DJ, Kan K, Bartels R, Satterfield BC, Riedy SM, et al. Naturalistic field study of the restart break in US commercial motor vehicle drivers: truck driving, sleep, and fatigue. *Accid Anal Prev*. 2016;93:55–64.
- Seelig AD, Jacobson IG, Smith B, Hooper TI, Boyko EJ, Gackstetter GD, et al. Sleep patterns before, during, and after deployment to Iraq and Afghanistan. *Sleep*. 2010;33(12):1615–22.
- Taylor MK, Hilton SM, Campbell JS, Beckerley SE, Shobe KK, Drummond SP, et al. Prevalence and mental health correlates of sleep disruption among military members serving in a combat zone. *Mil Med*. 2014;179(7):744–51.
- Troxel WM, Shih RA, Pedersen ER, Geyer L, Fisher MP, Griffin BA, Haas AC, Kurz J, Steinberg PS. Sleep in the military: promoting healthy sleep among US servicemembers. *Rand Health Q*. 2015;5(2). https://www.rand.org/content/dam/rand/pubs/research_reports/RR700/RR739/RAND_RR739.pdf. Accessed 19 May 2019.
- Taylor DJ, Pruiksma KE, Hale WJ, Kelly K, Maurer D, Peterson AL, et al. Prevalence, correlates, and predictors of insomnia in the US army prior to deployment. *Sleep*. 2016;39(10):1795–806.
- Mundry R, Nunn CL. Stepwise model fitting and statistical inference: turning noise into signal pollution. *Am Nat*. 2009;173(1):119–23.
- Baglioni C, Nanovska S, Regen W, Spiegelhalter K, Feige B, Nissen C, et al. Sleep and mental disorders: a meta-analysis of polysomnographic research. *Psychol Bull*. 2016;142(9):969–90.
- Ohayon MM. Prevalence of DSM-IV diagnostic criteria of insomnia: distinguishing insomnia related to mental disorders from sleep disorders. *J Psychiatr Res*. 1997;31(3):333–46.
- Mysliwiec V, McGraw L, Pierce R, Smith P, Trapp B, Roth BJ. Sleep disorders and associated medical comorbidities in active duty military personnel. *Sleep*. 2013;36(2):167–74.
- Cappuccio FP, Miller MA. Sleep and cardio-metabolic disease. *Curr Cardiol Rep*. 2017;19(11):110.
- Covassin N, Singh P. Sleep duration and cardiovascular disease risk: epidemiologic and experimental evidence. *Sleep Med Clin*. 2016;11(1):81–9.
- Pavloski P, Shelgikar AV. Treatment options for obstructive sleep apnea. *Neurol Clin Pract*. 2017;7(1):77–85.
- OTSG. Health of the Force report, 2015. <https://armymedicine.health.mil/Reports>. Accessed 20 May 2019.
- Thompson MA, Jones CB, Thornburg CJ. Sleep banking. *Mil Rev*. 2017:91–7.
- Lewis Shattuck N, Matsangas P, Moore J, Wegemann L. Prevalence of musculoskeletal symptoms, excessive daytime sleepiness, and fatigue in the crewmembers of a U.S. navy ship. *Mil Med*. 2016;181(7):655–62.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

